

Anti-hCG beta 5004 SP-1

Product overview

Catalog number	100004
Specificity	Antibody recognizes human chorionic gonadotropin and its free beta subunit
Description	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components.
Product buffer solution	0.9 % NaCl, 0.095 % NaN ₃ as a preservative
Shelf life and storage	36 months from manufacturing at 2–8 °C
Subclass	IgG ₁
Analyte description	Human chorionic gonadotropin (hCG) is a glycoprotein hormone produced in pregnancy by the developing embryo soon after conception and later by the syncytiotrophoblast (part of the placenta). Its role is to prevent the disintegration of the corpus luteum of the ovary and thereby maintain progesterone production that is critical for a pregnancy in humans. Early pregnancy testing, in general, is based on the detection of hCG. hCG is produced also by some tumors, but it is not known whether this production is a contributing cause or an effect of tumorigenesis.

Parameters tested on each lot

Product appearance	Liquid, may turn slightly opaque during storage
Product concentration	1.0 mg/ml (+/- 10%)
Immunoreactivity	80–120% compared to the reference sample in an FIA test
IEF Profile	6.5–7.6
Purity	≥ 95 %

Kinetic parameters

Association rate constant	hCG: 2.0×10^6 1/Ms and hCGβ: 5.0×10^5 1/Ms
Dissociation rate constant	hCG: 3.7×10^{-5} 1/s and hCGβ: 1.2×10^{-4} 1/s
Affinity constant	hCG: $K_A = 5.3 \times 10^{10}$ 1/M; $K_D = 1.9 \times 10^{-11}$ M (= 0.02 nM) hCGβ: $K_A = 4.3 \times 10^9$ 1/M; $K_D = 2.3 \times 10^{-10}$ M (= 0.23 nM)
Determination method	SPR (ProteOn XPR36)
Determination antigen	hCG, Scripps (Cat C0714, Lot 2430801); hCGβ, Scripps (Cat C0914, Lot 2310001)



Legal disclaimer

Cross-reactivities

LH ($K_D = 17$ nM)
Does not recognize hCG α , FSH, or TSH

Epitope

Beta-2 as described in Berger et al. (2013). The antibody recognizes both intact hCG and free β subunit.

Pair recommendations

		DETECTION										
		hCG beta								alpha subunit		
		5004	5006	5008	5009	5011	5012 free β	5014	5016	5501	5503	6601
CAPTURE	5004	-	-	-	+	+	-	+	-	+	+	+
	5006	-	-	-	-	-	-	+	-	+	+	+
	5008	-	-	-	+	-	-	+	-	+	+	+
	5009	+	+	+	-	-	-	+	+	-	-	+
	5011	+	+	+	-	-	-	+	+	-	-	+
	5012 free β	+	+	+	-	-	-	+	+	-	-	-
	5014	+	+	+	+	+	-	-	+	+	+	+
	5016	-	-	-	-	-	-	+	-	+	+	+

Following pairs are especially recommended for free hCG beta assays:
CLIA: 5012 (capture) – 5004 (detection) and 5012 – 5008

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Platforms tested

FIA, CLIA

Antigens tested

Native hCG antigens, Lee Biosolutions 189-10 and 189-11
Native β -hCG antigen, Lee Biosolutions 325-11

Product stability

TEMPERATURE, TIME RESULT
Not Determined (N/D) -

Miscellaneous

-

References

Berger, P., Paus, E., Hemken, P.M., Sturgeon, C., Stewart, W.W., Skinner, J.P., Harwick, L.C., Saldana, S.C., Ramsay, C.S., Rupprecht, K.R., Olsen, K.H., Bidart, J.M. and Stenman, U.H. (2013) Candidate epitopes for measurement of hCG and related molecules: the second ISOBM TD-7 workshop. Tumor Biol., 34: 4033-4057.



Legal disclaimer